



# Dr Georges MOUTON MD

Functional Medicine

## QUOTE GM #38

06/12/2018

Titre

Créé le

### LE MICROBIOTE INTESTINAL JOUE AUSSI UN RÔLE DANS L'HYPERTENSION

Front Cell Infect Microbiol. 2017 Aug 24;7:381. doi: 10.3389/fcimb.2017.00381. eCollection 2017.

#### Alterations of the Gut Microbiome in Hypertension.

Yan Q<sup>1,2</sup>, Gu Y<sup>3</sup>, Li X<sup>4</sup>, Yang W<sup>5</sup>, Jia L<sup>1</sup>, Chen C<sup>1</sup>, Han X<sup>1</sup>, Huang Y<sup>1</sup>, Zhao L<sup>1</sup>, Li P<sup>3</sup>, Fang Z<sup>3</sup>, Zhou J<sup>3</sup>, Guan X<sup>5</sup>, Ding Y<sup>6</sup>, Wang S<sup>7</sup>, Khan M<sup>8</sup>, Xin Y<sup>9</sup>, Li S<sup>3</sup>, Ma Y<sup>1</sup>.

Author information

#### Abstract

**Introduction:** Human gut microbiota is believed to be directly or indirectly involved in cardiovascular diseases and hypertension. However, the identification and functional status of the hypertension-related gut microbe(s) have not yet been surveyed in a comprehensive manner. **Methods:** Here we characterized the gut microbiome in hypertension status by comparing fecal samples of 60 patients with primary hypertension and 60 gender-, age-, and body weight-matched healthy controls based on whole-metagenome shotgun sequencing. **Results:** Hypertension implicated a remarkable gut dysbiosis with significant reduction in within-sample diversity and shift in microbial composition. Metagenome-wide association study (MGWAS) revealed 53,953 microbial genes that differ in distribution between the patients and healthy controls (false discovery rate, 0.05) and can be grouped into 68 clusters representing bacterial species. Opportunistic pathogenic taxa, such as, *Klebsiella* spp., *Streptococcus* spp., and *Parabacteroides merdae* were frequently distributed in hypertensive gut microbiome, whereas the short-chain fatty acid producer, such as, *Roseburia* spp. and *Faecalibacterium prausnitzii*, were higher in controls. The number of hypertension-associated species also showed stronger correlation to the severity of disease. Functionally, the hypertensive gut microbiome exhibited higher membrane transport, lipopolysaccharide biosynthesis and steroid degradation, while in controls the metabolism of amino acid, cofactors and vitamins was found to be higher. We further provided the microbial markers for disease discrimination and achieved an area under the receiver operator characteristic curve (AUC) of 0.78, demonstrating the potential of gut microbiota in prediction of hypertension. **Conclusion:** These findings represent specific alterations in microbial diversity, genes, species and functions of the hypertensive gut microbiome. Further studies on the causality relationship between hypertension and gut microbiota will offer new prospects for treating and preventing the hypertension and its associated diseases.

**KEYWORDS:** gut microbiome; hypertension; metagenome-wide association study; microbial dysbiosis

PMID: 28884091 PMCID: PMC5573791 DOI: 10.3389/fcimb.2017.00381

**Introduction :** *Il semble que le microbiote intestinal humain soit directement ou indirectement impliqué dans les maladies cardiovasculaires et dans l'hypertension.*

**Résultats :** *L'hypertension impliquait une dysbiose intestinale remarquable avec une réduction significative de la diversité microbienne au sein de l'échantillon et une modification de cette composition microbienne.*

**Conclusion :** *Ces résultats reflètent des altérations spécifiques de la diversité microbienne, des gènes, des espèces et des fonctions du microbiome intestinal présent en cas d'hypertension."*